

REFERENCE CHART: HOLDING TIME, BOTTLE TYPE, PRESERVATION, MINIMUM VOLUME

Note: Multiple analyses may be run from the same container, provided that the bottle type, volume and preservation are appropriate.

Analytical Parameter	Maximum Holding Time	Required Container Type	Required Preservative	Minimum Amount	
Alkalinity (Bicarb, Carb, Hyd, and Tot); SM 2320 B	14 days	Plastic / Glass	0-6°C	100	mL
Ammonia (NH3 as N); SM 4500NH3 B/C or B/G	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	200	mL
Anions by EPA 300.0: Chloride (Cl ⁻); Fluoride (F ⁻);	28 days	Plastic / Glass	0-6° C (no temp. req. for	100	mL
Nitrate (as N); Sulfate (SO ₄ ²⁻)	48 hours (Nitrate)	. ideas, ciaes	Chloride or Fluoride)		
Asbestos (TEM)	48 hours	Plastic / Glass	0-6°C	1000	mL
Bioassay (Acute Tox) % survival	36 hours	Plastic cubitainer	0-6°C	1-5	gal
Biochemical Oxygen Demand (BOD); SM 5210 B	48 hours	Plastic / Glass	0-6°C	500	
Chromium, Hexavalent (CrVI); SM 3500Cr B	24 hours		0-6°C	200	mL ml
		Plastic / Glass	0-6 C	200	mL
Chlorine, residual; SM 4500Cl B or G	15 minutes	Field Measurement	-	-	
Chemical Oxygen Demand (COD); SM 5220 D or EPA 410.4	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	50	mL
Chlorophyll A (Algal Biomass) / Pheophytin; SM 10200 H	48 Hours to filter 28 days once filtered	Amber Plastic	0-6°C, unfiltered -20°C, filtered	1	L
	8 hrs -	Plastic (sterile)	$Na_2S_2O_3 + 0-10^{\circ}C;$	100	mL
Coliform, Total / Fecal; SM 9221 B/E	wastewater/stormwater	, ,	No regulatory temp. req.		
·, · · · · · · · · · · · · · · · · · ·	30 hrs - drinking water		for drinking water)		
Coliform, Total / E.Coli; SM 9223 B	30 hrs – drinking water	Plastic (sterile)	Na ₂ S ₂ O ₃ + 0-10°C; No	100	mL
(Present/Absent <u>or</u> Quantitray)	l come annual and	r ideae (eterne)	regulatory temp. req. for DW		
Color; SM 2120 B	48 hours	Plastic / Glass	0-6°C	100	mL
Conductivity (EC/SC); SM 2510 B	28 days	Plastic / Glass	0-6°C	100	mL
Conductivity (EG/CG), GW 2310 B	14 days			100	mL
Cyanide Total; SM 4500CN C/E	14 days	Amber plastic with NaOH pH >10 + 0-6°C Pre-treatment for Total Cyanide samples with chlorine or NO ₃ /NO ₂ should be done prior to preservation. Kits can be provided upon request.			IIIL
Cyanide WAD; SM 4500CN I/E	14 days	Amber plastic	NaOH pH >10 + 0-6°C	100	mL
Dissolved Organic Carbon (DOC); SM 5310 B	Filter within 48 hours 28 days	125 mL Amber glass	Filter then add HCl to pH <2 + 0-6°C	50	mL
Dissolved Oxygen (DO); SM 45000 G	15 minutes	Field Measurement or Glass bottle/no headspace Collect in duplicate	0-6°C	500	mL
Enterococcus by Enterolert	8 hours	Plastic (sterile)	Na ₂ S ₂ O ₃ + 0-10°C	100	mL
Heterotrophic Plate Count (HPC/SPC); SM 9215E (SimPlate)	8 hours	Plastic (sterile)	Na ₂ S ₂ O ₃ + 0-10°C	100	mL
Hardness (by titration); SM 2340 C	6 months	Plastic / Glass	HNO₃ pH <2	250	mL
,	Filter within 15	Plastic / Glass {15}	Filter then add HNO ₃	100	mL
Metals, Dissolved {2}; EPA 200.8 Metals, Total; EPA 200.8	minutes {2}, 6 months		HNO ₃ pH <2	100	
Metals, Total; EPA 6010/6020	6 months	Plastic / Glass {15} Solids in Glass SJ	None	1	mL 8oz
Mercury (Hg) EPA 245.1	28 days	Plastic / Glass	HNO ₃ pH <2	100	mL
Mercury (Hg) EPA 7471		Solids in Glass SJ	0-6°C	1	8oz
Mercury, Total; EPA 1631 (Aqueous)	48 hours to preserve 90 days once preserved	Glass, Dbl Bagged (not appropriate for samples with solids TSS>200 mg/L)	HCI	100	mL
Mercury, Dissolved {2}; EPA 1631	Filter within 24 hours, Preserve within 48 hours 90 days once preserved	Glass, Dbl Bagged	Filter in lab, then HCl	100	mL
Mercury, Methyl; EPA 1630 (Aqueous)	Preserve {8} within 48 hrs 6 months preserved	Glass, Amber Dbl Bagged	Dark and cool + (HCl or H ₂ SO ₄ {8})	100	mL
Mercury, Methyl; EPA 1630 (Sludge or Solids)	6 months frozen	Solids in Amber Poly SJ	Solids to be Frozen	1	8oz
Mercury, Dissolved {2} Methyl; EPA 1630	Filter & preserve within 48 hrs	Glass, Amber Dbl Bagged	Filter in lab, dark and cool + (HCl or H ₂ SO ₄ {8})	100	mL
Nitrate (as NI): FDA 200 0	6 months once preserved	Plantia / Class	0.6%	100	mal
Nitrate (as N); EPA 300.0	48 hours	Plastic / Glass	0-6°C	100	mL ml
Nitrite (as N); SM 4500NO2 B	48 hours	Plastic / Glass	0-6°C	100	mL
Nitrate+Nitrite as N (NO ₃ +NO ₂ -N) EPA 353.2, or SM NO3 F	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	100	mL
	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	200	mL
Nitrogen, Total Kjeldahl (TKN); SM 4500Norg + SM 4500NH3 B					mL
	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	200	IIIL
4500NH3 B	6 hours (recommended)	Plastic / Glass Glass only	H ₂ SO ₄ pH <2 +0-6°C 0-6°C	1000	mL
4500NH3 B Nitrogen, Total Organic (TON) TKN-NH ₃ = TON calc Odor; SM 2150 B	6 hours (recommended) / 24 hour regulatory	Glass only	0-6°C	1000	mL
4500NH3 B Nitrogen, Total Organic (TON) TKN-NH ₃ = TON calc	6 hours (recommended)				



Analytical Parameter	Maximum Holding Time	Required Container Type	Required Preservative	Minimum Amount	
Polynuclear Aromatic Hydrocarbons (PAH's)	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C {5}	1	L
EPA 625.1/8270	14 days, Sludge or Solid	triplicate(6). Solids in Glass SJ	0-6°C	1	8oz
Organophosphorus Pesticides	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C {5}	1	L
EPA 625.1 (formerly EPA 614) /8141	14 days, Sludge or Solid	triplicate(6). Solids in Glass SJ	0-6°C	1	8oz
Semi-volatile Organics	7 days, Aqueous	Glass Amber Liter - Collect in triplicate(6). Solids in Glass SJ	0-6°C {5}	1 1	L
EPA 625.1/8270	14 days, Sludge or Solid 7 days /3 days {9}	Glass Amber Liter - Collect in	0-6°C 0-6°C {5}	<u>1</u>	8oz L
Pyrethroid Pesticides	7 days 73 days {9}	triplicate(6).	0-6 C {5}		L
GCMS-NCI-SIM/ EPA 8270/625.1	14 days, soil or sediment	Solids in Amber Glass SJ	Solids to be Frozen	1	8oz
Carbamate Pesticides	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C	1	L
EPA 632/8321	14 days, Sludge or Solid	triplicate(6). Solids in Glass SJ	0-6°C	1	8oz
Volatile Organics	14 days, preserved	Glass VOA vial {1} (sample	HCl pH <2 + 0-6°C {4}	40	mL
EPA 524.2 {4}	l radye, preserved	in triplicate)	1101 pit 2 1 0 0 0 (1)		
	14 days except:	Min. of four 40 mL VOA Vials.	0-6°C		
Volatile Organics	3 days unpreserved for	6-pack preferred; Two Pres w/	(see holding time for	One 40	
EPA 624.1 {3}	Acrolein {12}	HCl, Two Unpreserved, and	additional preservation)		ontainer
(Includes Acrolein and Acrylonitrile)	7 days if only unpreserved	Two pH 4-5 {1}	μ,	type de	escribed
	received {11}.	THMS only—3 HCI VOAs			
	14 days (aqueous)	Three Clear Glass VOA Vials{1}	HCl pH <2 + 0-6°C	40	mL
Volatile Organics; EPA 8260 {3}	(aqueeus)	Solids in separate glass jar	1161 pri 12 1 0 0 0		
1 olamo olganios, 21 / 1 o2 o (e)	14 days Sludge/solid	filled to top	0-6°C	1	8oz
TPH Diesel/Motor Oil; EPA 8015	7 davs	Glass Amber Liter {6}	0-6°C	1	L
Tributyltin (TBT)	7 days (recommended)	Glass Amber Liter {6}	HCl pH <2 + 0-6°C	1	L
TPH Gas/ BTEX/ MTBE; EPA 8260	14 days preserved	Glass VOA vial {1}	HCl pH <2 + 0-6°C	40	mL
Dioxin; EPA 1613	1 year	Glass Amber Liter {6}	0-6°C {5}	1	L
Perchlorate; EPA 314.0	28 days	Plastic/Glass	0-6°C	100	mL
pH; SM 4500 H+ B	15 minutes	Plastic / Glass	0-6°C	100	mL
Phenols, EPA 420.4	28 days	Glass Amber 250 mL	H ₂ SO ₄ pH <2 +0-6°C	200	mL
Phosphate, Ortho (as P); SM 4500P E	48 hours	Plastic / Glass	0-6°C	100	mL
,	15 minutes to filter /	Plastic / Glass	0-6°C	100	mL
Phosphate, Ortho, Dissolved (as P); SM 4500P E	48 hrs once filtered				
Phosphorus, Total (as P); SM 4500P B/F or B/E	28 days	Plastic / Glass	H ₂ SO ₄ pH <2+0-6°C	100	mL
Solids, Settleable (SS); SM 2540 F	48 hours	Plastic / Glass	0-6°C	1000	mL
Solids, Total (mg/L or %); SM 2540 B, or SM 2540 G	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Total Dissolved (TDS); SM 2540 C	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Total Suspended (TSS); SM 2540 D	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Volatile Suspended (VSS); SM 2540 E	7 days	Plastic / Glass	0-6°C	200	mL
Sulfide, Dissolved; SM 4500 S2- B/D	7 days	Glass 250mL (no headspace)	NaOH + AICI ₃ +0-6°C {13}	250	mL
Sulfide, Total; SM 4500 S2- C/D	7 days	Glass (preferred)/plastic	NaOH + ZnAC	250	mL
		(no headspace)	pH >9 +0-6°C		
Sulfite (SO ₃)	7 days	Glass Amber 500 mL	EDTA + 0-6°C	250	mL
Surfactants (MBAS); SM 5540 C	48 hours	Plastic / Glass	0-6°C	250	mL
Suspended Sediment Conc. (SSC); ASTM D3977	7 days	Plastic / Glass	0-6°C {7}	100	mL
Total Organic Carbon (TOC); SM 5310 B	28 days	Amber Glass VOA (3 vials)	HCl pH <2 + 0-6°C	40	mL
Turbidity; SM 2130 B	48 hours	Plastic / Glass	0-6°C	100	mL
Ultraviolet Absorption (UVA at 254nm); SM 5910 B	48 hours	Glass Amber 125mL	0-6°C	125	mL

- {1} Volatile organic samples need to be filled in multiple VOA vials without air bubbles/headspace (≤6 mm in size).
- (2) Dissolved metals require field or lab filtration through 0.45-micron filter prior to preservation. 40 CFR 136.3 requires filtration within 15 minutes.
- (3) Volatile organic methods EPA 624.1 and 8260 require dechlorination using Sodium Thiosulfate (Na₂S₂O₃) at time of sampling if chlorine is present. (Thiosulfate dechlorination bottles are available at the laboratory upon request.) Dechlorination is to occur before transferring to the appropriate VOA.
- (4) Volatile organic method EPA 524.2 requires dechlorination using ascorbic acid at time of sampling if chlorine is present. Ascorbic acid dechlorination kits are available at the laboratory upon request. If analyzing for THMs only, Sodium Thiosulfate (Na₂S₂O₃) may be used, and acidification can be omitted. Otherwise, dechlorinate with Ascorbic acid, then preserve with HCl. If the sample foams vigorously upon addition of HCl, discard and collect unpreserved dechlorinated sample, and notify the laboratory as the samples must be analyzed within 24 hours of collection if they are to be analyzed for any compounds other than THMs. Method 524.2 requires a travel/trip blank with each sample set collected.
- (5) If sampling from a chlorinated location, add 80 mg/L Sodium Thiosulfate (Na₂S₂O₃) per liter and mix well. Any method suitable for field use may be employed to test for residual chlorine (Reference 16). Add more Sodium Thiosulfate if 80 mg/L is insufficient but do not add excess Sodium Thiosulfate.
- (6) Semi Volatile Extractable Organics in Amber Liters (AL) should be collected in enough bottles to ensure the lab can perform method-required Matrix Spike/Spike Duplicate (MS/MSD) analyses. While 3 AL per method is recommended, when collecting AL's for multiple methods, the number of AL's per method can be reduced to 2 per method. Please contact your Project Manager if in doubt about number of AL's per sample or method.
- {7} Suspended Sediment Concentration (SSC) requires its own container and the entire contents are used for the analysis.
- (8) Preserve with HCl if less than 10 ppT Salinity, or preserve with H2SO4 if greater than 10 ppT Salinity.
- (9) Cyhalothrin in water has a 3-day hold time in reagent water per USGS study. Permethrin in water has a 3-day hold time per Storage Stability Study by CA Dept. of Food and Agriculture.
- (10) 1L bottle required for lowest available MDL or RL <5 mg/L; 500mL bottle required for RL of 5 mg/L; 250mL bottle recommended for samples with expected concentrations of 20 mg/L or higher.
- {11} If unpreserved, must be analyzed within 7 days of sampling.
- {12} Hold time can be extended to 14 days if preserved to pH 4-5 at time of collection.
- (13) Return sample to laboratory the same day as collection for pH verification, decanting, and further preservation.
- (14) Samples with visible solids may only require 100mL. Samples with no or low visible solids should be collected in 1000mL containers.
- {15} Glass not acceptable for boron.



(Rev 09/14/2023)

