

REFERENCE CHART: HOLDING TIME, BOTTLE TYPE, PRESERVATION, MINIMUM VOLUME Note: Multiple analyses may be run from the same container, provided that the bottle type, volume and preservation are appropriate.

Analytical Parameter	Maximum Holding Time	Required Container Type	Required Preservative	Minimum Amount	
Alkalinity (Bicarb, Carb, Hyd, and Tot); SM 2320 B	14 days	Plastic / Glass	0-6°C	100	mL
Ammonia (NH3 as N); SM 4500NH3 B/C or B/G	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	200	mL
Anions by EPA 300.0: Chloride (Cl ⁻); Fluoride (F ⁻);	28 days	Plastic / Glass	0-6° C (no temp. req. for	100	mL
Nitrate (as N); Sulfate (SO ₄ ²⁻)	48 hours (Nitrate)		Chloride or Fluoride)		
Asbestos (TEM)	48 hours	Plastic / Glass	0-6°C	1000	mL
Bioassay (Acute Tox) % survival	36 hours	Plastic cubitainer	0-6°C	1-5	
Biochemical Oxygen Demand (BOD); SM 5210 B	48 hours	Plastic / Glass	0-6°C	500	mL
Chromium, Hexavalent (CrVI); SM 3500Cr B or EPA	24 hours	Plastic / Glass	0-6°C	200	mL
7196			0-0 0	200	
Chlorine, residual; SM 4500Cl B or G	15 minutes	Field Measurement	-	-	-
Chemical Oxygen Demand (COD); SM 5220 D	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	50	mL
Chlorophyll A (Algal Biomass) / Pheophytin; SM 10200 H	48 Hours to filter 28 days once filtered	Amber Plastic	0-6°C, unfiltered -20°C, filtered	1	L
1020011	8 hrs –	Plastic (sterile)	$Na_2S_2O_3 + 0-10^{\circ}C;$	100	mL
Coliform Total / Fecal: SM 9221 B/F	wastewater/stormwater		No regulatory temp. req.	100	
Coliform, Total / Fecal; SM 9221 B/E	30 hrs – drinking water				
	-		for drinking water)	100	
Coliform, Total / E.Coli; SM 9223 B (Present/Absent <u>or</u> Quantitray)	30 hrs – drinking water	Plastic (sterile)	Na ₂ S ₂ O ₃ + 0-10°C; No regulatory temp. req. for DW	100	mL
Color; SM 2120 B	48 hours	Plastic / Glass	0-6°C	100	mL
Conductivity (EC/SC); SM 2510 B	28 days	Plastic / Glass	0-6°C	100	mL
• • •	14 days	Amber plastic with NaOH pH >		100	mL
Cyanide Total; SM 4500CN C/E	,	Pre-treatment for Total Cyanide samples with chlorine or NO ₃ /NO ₂ should be done prior to preservation. Kits can be provided upon request.			
Cyanide WAD; SM 4500CN I/E	14 days	Amber plastic	NaOH pH >10 + 0-6°C	100	mL
Dissolved Organic Carbon (DOC); SM 5310 B	Filter within 48 hours	125 mL Amber glass	Filter then add HCl to	50	mL
Dissolved Organic Carbon (DOC), Sivi 5310 B	28 days	Field Measurement on Olana	pH <2 + 0-6°C	500	mal
Dissolved Oxygen (DO); SM 4500O G	15 minutes	Field Measurement or Glass bottle/no headspace Collect in duplicate	0-6°C	500	mL
Enterococcus by Enterolert	8 hours	Plastic (sterile)	Na ₂ S ₂ O ₃ + 0-10°C	100	mL
Heterotrophic Plate Count (HPC/SPC); SM 9215E (SimPlate)	8 hours	Plastic (sterile)	$Na_2S_2O_3 + 0-10^{\circ}C$	100	mL
Hardness (by titration); SM 2340 C	6 months	Plastic / Glass	HNO₃ pH <2	250	mL
Metals, Dissolved {2}; EPA 200.8	Filter within 15	Plastic / Glass {15}	Filter then add HNO ₃	100	mL
Matala Tatali EDA 200.0	minutes {2}, 6 months	Directio / Class (15)		100	
Metals, Total; EPA 200.8 Metals, Total; EPA 6010/6020	6 months	Plastic / Glass {15} Solids in Glass SJ	HNO ₃ pH <2 None	100 1	mL 8oz
Mercury (Hg) EPA 245.1	28 days	Plastic / Glass	HNO ₃ pH <2	100	mL
Mercury (Hg) EPA 7471	-	Solids in Glass SJ	0-6°C	1	8oz
Mercury, Total; EPA 1631 (Aqueous)	48 hours to preserve 90 days once preserved	Glass, Dbl Bagged (not appropriate for samples with solids TSS>200 mg/L)	HCI	100	mL
	Filter within 24 hours.	Glass, Dbl Bagged	Filter in lab. then HCl	100	mL
Mercury, Dissolved {2}; EPA 1631	Preserve within 48 hours 90 days once preserved	Class, Dbi Dagged		100	
Mercury, Methyl; EPA 1630 (Aqueous)	Preserve {8} within 48 hrs	Glass, Amber Dbl Bagged	Dark and cool + (HCl or	100	mL
moroary, morry, Er A 1000 (Aqueous)	6 months preserved	Class, Alliper Doi Dayyeu	H_2SO_4 {8})	100	
Mercuny Methyl: EDA 1630 (Sludge or Solide)	6 months frozen	Solids in Amber Poly S I		1	8oz
Mercury, Methyl; EPA 1630 (Sludge or Solids)		Solids in Amber Poly SJ Glass, Amber Dbl Bagged	Solids to be Frozen Filter in lab, dark and cool	100	
Mercury, Dissolved {2} Methyl; EPA 1630	Filter & preserve within 48 hrs	Glass, Amber Dbi Bagged	+ (HCl or H_2SO_4 {8})	100	mL
	6 months once preserved	Dia the / Ole as	0.0%0	400	
Nitrate (as N); EPA 300.0	48 hours	Plastic / Glass	0-6°C	100	mL
Nitrite (as N); SM 4500NO2 B	48 hours	Plastic / Glass	0-6°C	100	mL
Nitrate+Nitrite as N (NO ₃ +NO ₂ -N) EPA 353.2, or SM 4500-NO3 F	28 days	Plastic / Glass	H₂SO₄ pH <2 +0-6°C	100	mL
Nitrogen, Total Kjeldahl (TKN); SM 4500Norg + SM 4500NH3 B	28 days	Plastic / Glass	H₂SO₄ pH <2 +0-6°C	200	mL
Nitrogen, Total Organic (TON) TKN-NH ₃ = TON calc	28 days	Plastic / Glass	H ₂ SO ₄ pH <2 +0-6°C	200	mL
Odor; SM 2150 B	6 hours (recommended) / 24 hour regulatory	Glass only	0-6°C	1000	mL
Oil & Grease, Total and/or Hydrocarbons EPA 1664	28 days	Glass only-in duplicate	HCI or H ₂ SO₄ pH<2+0-6°C	500	mL{10}
Chlorinated Pesticides & PCBs, EPA 625.1	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C {5}	1	
(formerly EPA 608) /8081/8082	14 days, Aqueous 14 days, Sludge or Solid	triplicate{6}. Solids in Glass SJ	0-6°C	-	L 8oz
(10111011) EFA 000/10001/0002	14 days, sludge of solid	anpiroate of. Dolids in Glass 30	0-0 C	1	002



(Rev 10/16/2024)

CA-ELAP Certification 1664

Analytical Parameter	Maximum Holding Time	Required Container Type	Required Preservative	Minimum Amount	
Organophosphorus Pesticides	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C {5}	1	L
EPA 625.1 (formerly EPA 614) /8141	14 days, Sludge or Solid	triplicate{6}. Solids in Glass SJ	0-6°C	1	8oz
Semi-volatile Organics/	7 days, Aqueous	Glass Amber Liter - Collect in	0-6°C {5}	1	L
PAHs by EPA 625.1/8270	14 days, Sludge or Solid	triplicate{6}. Solids in Glass SJ	0-6°C	1	8oz
Pyrethroid Pesticides	7 days /3 days {9}	Glass Amber Liter - Collect in	0-6°C {5}	1	L
GCMS-NCI-SIM/ EPA 8270/625.1	14 days sail or addiment	triplicate{6}.	Colida ta ha Frazan	1	0.07
Carbamate Pesticides	14 days, soil or sediment 7 days, Aqueous	Solids in Amber Glass SJ Glass Amber Liter - Collect in	Solids to be Frozen 0-6°C	<u>1</u> 1	8oz
EPA 632/8321	14 days, Aqueous 14 days, Sludge or Solid	triplicate{6}. Solids in Glass SJ	0-6°C	1	L 8oz
		,		40	-
Volatile Organics	14 days, preserved	Glass VOA vial {1} (sample	HCl pH <2 + 0-6°C {4}	40	mL
EPA 524.2 {4}	14 days except:	in triplicate) Min. of four 40 mL VOA Vials.	0-6°C		
Malatila Ormaniaa	3 days unpreserved for	6-pack preferred; Two Pres w/	(see holding time for	One 40)mL of
Volatile Organics	Acrolein {12}	HCI, Two Unpreserved, and	additional preservation)	each container	
EPA 624.1 {3}	7 days if only unpreserved	Two pH 4-5 {1}	additional preservation)	type de	escribed
(Includes Acrolein and Acrylonitrile)	received {11}.				
		THMS only—3 HCI VOAs Three Clear Glass VOA Vials{1}	HCl pH <2 + 0-6°C	40	
Volatile Organics; EPA 8260 {3}	14 days (aqueous)	Solids in separate glass jar	HCI pH <2 + 0-6 C	40	mL
	14 days Sludge/solid	filled to top	0-6°C	1	8oz
TPH Diesel/Motor Oil; EPA 8015	7 days	Glass Amber Liter {6}	0-6°C	1	1
Tributyltin (TBT)	7 days 7 days (recommended)	Glass Amber Liter {6}	HCI pH <2 + 0-6°C	1	
TPH Gas/ BTEX/ MTBE; EPA 8260	14 days preserved	Glass VOA vial {1}	HCl pH <2 + 0-6°C	40	mL
Dioxin: EPA 1613	1 year	Glass Amber Liter {6}	0-6°C {5}	<u>+0</u> 1	1
Perchlorate; EPA 314.0	28 days	Plastic/Glass	0-6°C	100	mL
pH; SM 4500 H+ B	15 minutes	Plastic / Glass	0-6°C	100	mL
Phenols, EPA 420.4	28 days	Glass Amber 250 mL	H₂SO₄ pH <2 +0-6°C	200	mL
Phosphate, Ortho (as P); SM 4500P E	48 hours	Plastic / Glass	0-6°C	100	mL
	15 minutes to filter /	Plastic / Glass	0-6°C	100	mL
Phosphate, Ortho, Dissolved (as P); SM 4500P E	48 hrs once filtered	Flastic / Glass	0-0 C	100	···· L
Phosphorus, Total (as P); SM 4500P B/F or B/E	28 days	Plastic / Glass	H₂SO₄ pH <2+0-6°C	100	mL
Solids, Settleable (SS); SM 2540 F	48 hours	Plastic / Glass	0-6°C	1000	mL
Solids, Total (mg/L or %); SM 2540 B, or SM 2540 G	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Total Dissolved (TDS); SM 2540 C	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Total Suspended (TSS); SM 2540 D	7 days	Plastic / Glass	0-6°C	1000	mL{14}
Solids, Volatile Suspended (VSS); SM 2540 E	7 days	Plastic / Glass	0-6°C	200	mL
Sulfide, Dissolved; SM 4500 S2- B/D	7 days	Glass 250mL (no headspace)	NaOH + AICI ₃ +0-6°C {13}	250	mL
Sulfide, Total; SM 4500 S2- C/D	7 days	Glass (preferred)/plastic	NaOH + ZnAC	250	mL
	7 days	(no headspace)	pH >9 +0-6°C	200	
Sulfite (SO ₃)	7 days	Glass Amber 500 mL	EDTA + 0-6°C	250	mL
Surfactants (MBAS); SM 5540 C	48 hours	Plastic / Glass	0-6°C	250	mL
Suspended Sediment Concentration (SSC)	7 days (recommended)				
ASTM D3977-97 B - Filtration	120 days max (per USGS)	Plastic / Glass	0-6°C {7}	100	mL
Total Organic Carbon (TOC); SM 5310 B	28 days	Amber Glass VOA (3 vials)	HCl pH <2 + 0-6°C	40	mL
Turbidity; SM 2130 B	48 hours	Plastic / Glass	0-6°C	100	mL
Ultraviolet Absorption (UVA at 254nm); SM 5910 B	48 hours	Glass Amber 125mL	0-6°C	125	mL

{1} Volatile organic samples need to be filled in multiple VOA vials without air bubbles/headspace (≤6 mm in size).

{2} Dissolved metals require field or lab filtration through 0.45-micron filter prior to preservation. 40 CFR 136.3 requires filtration within 15 minutes.

(3) Volatile organic methods EPA 624.1 and 8260 require dechlorination using Sodium Thiosulfate (Na2S2O3) at time of sampling if chlorine is present. (Thiosulfate dechlorination bottles are available at the laboratory upon request.) Dechlorination is to occur before transferring to the appropriate VOA. 4 Volatile organic method EPA 524.2 requires dechlorination using ascorbic acid at time of sampling if chlorine is present. Ascorbic acid dechlorination kits are available at the laboratory upon request. If analyzing for THMs only, Sodium Thiosulfate (Na₂S₂O₃) may be used, and acidification can be omitted. Otherwise, dechlorinate with Ascorbic acid, then preserve with HCI. If the sample foams vigorously upon addition of HCI, discard and collect unpreserved dechlorinated sample, and notify the laboratory as the samples must be analyzed within 24 hours of collection if they are to be analyzed for any compounds other than THMs. Method 524.2 requires a travel/trip blank with each sample set collected.

[5] If sampling from a chlorinated location, add 80 mg/L Sodium Thiosulfate (Na2S2O3) per liter and mix well. Any method suitable for field use may be employed to test for residual chlorine (Reference 16). Add more Sodium Thiosulfate if 80 mg/L is insufficient but do not add excess Sodium Thiosulfate. (6) Semi Volatile Extractable Organics in Amber Liters (AL) should be collected in enough bottles to ensure the lab can perform method-required Matrix

Spike/Spike Duplicate (MS/MSD) analyses. While 3 AL per method is recommended, when collecting AL's for multiple methods, the number of AL's per method can be reduced to 2 per method. Please contact your Project Manager if in doubt about number of AL's per sample or method.

[7] Suspended Sediment Concentration (SSC) requires its own container and the entire contents are used for the analysis.

(8) Preserve with HCl if less than 10 ppT Salinity, or preserve with H₂SO₄ if greater than 10 ppT Salinity.

(9) Cyhalothrin in water has a 3-day hold time in reagent water per USGS study. Permethrin in water has a 3-day hold time per Storage Stability Study by CA Dept. of Food and Agriculture.

(10) 1L bottle required for lowest available MDL or RL <5 mg/L; 500mL bottle required for RL of 5 mg/L; 250mL bottle recommended for samples with expected concentrations of 20 mg/L or higher.

{11} If unpreserved, must be analyzed within 7 days of sampling.

(12) Hold time can be extended to 14 days if preserved to pH 4-5 at time of collection.

(13) Return sample to laboratory the same day as collection for pH verification, decanting, and further preservation.

(14) Samples with visible solids may only require 100mL. Samples with no or low visible solids should be collected in 1000mL containers.

{15} Glass not acceptable for boron.



(Rev 10/16/2024)

